


PALEOZOIC PETROLEUM SYSTEMS OF THE WEST SIBERIAN BASIN - WHAT IS THE EVIDENCE?

Enver ABLYA¹, Ekaterina BORDY¹, Evgeniya KODLAEVA¹, Tatiana KORNEVA¹, Dmitryi NADEZHKIN¹, Pim VAN BERGEN², Ron HOFLAND² and Mark SUGDEN²

¹Moscow State University, Department of Petroleum Geology, Vorobjevy Gory, Moscow, 119992, RUSSIA

²Shell International Exploration and Production B.V. Kessler Park 1, 2288 GS Rijswijk, NL

Mesozoic petroleum systems in the West Siberian Basin have been studied extensively and are relatively well understood (Goncharov 1988, Kontorovich et al. 1996, Rudkevich et al. 1988, Peters et al. 1994). The two main source systems are the Upper Jurassic marine Bazhenov suite (J₃) and the Lower-Middle Jurassic more terrigenously influenced Togur/Tyumen suites (J₁-J₂). Most of the oils known to be generated by these sources are produced from Mesozoic reservoirs. However, there is also production from Paleozoic strata. The petroleum systems related to these latter reservoirs are much less understood, in particular with respect to source, and form the basis of this study. The oils studied here are from Paleozoic reservoirs located in the western part of the basin around Uray and the south-eastern area around Tomsk.



Samples

South-eastern area (Red)

1. Eley-Igayskoe, 2, 3808 m
2. Kalinovoe, 6, 2970-2980 m
3. Maloichskoe, 2, 2842-2849 m
4. Nizhne-Tabaganskoe, 4, 3068-3080 m
5. Yzhno-Tabaganskoe, 130, 2981-3012 m
6. Severo-Ostaninskoe, 7, 2794-2810 m
7. Urmanskoe, 2, 3091-3103 m
8. Verhnetarskoe, 3, 2692-2704 m

Western area (Yellow)

9. Rogozhnikovskoe, 729, 2568-2607 m
10. Yahlinskoe, 851, 4025-4070 m
11. Palyanovskoe, 51, 2219-2422 m
12. Ubinskoe, 328, 1879-1913 m
13. Hanti-Mansiyskoe, 5, 3118-3152 m

Figure 1. Distribution of $\alpha\beta$ C₂₇₋₂₉ steranes

The oils are produced mainly from Middle-Upper Devonian horizons with one sample from a Lower Carboniferous horizon. In the literature these reservoirs are described as residual soil structures of Paleozoic rocks, or as Paleozoic complexes. They are often porous, karst-type reservoirs in organic rich limestone, occasionally with admixtures of dolomite, or with layers of clayey limestone.

Distinct sources were recognized based on the general molecular composition (e.g. *n*-alkanes, isoprenoids, naphthalenes) of whole oil, saturate and aromatic fractions, in combination with detailed biomarker evaluations. Comparisons with other oils from the West Siberian Basin showed that the two main contributions were from the two well known Jurassic sources. A characteristic even over odd predominance of the C₂₂, C₂₄ and C₂₆ *n*-alkanes, a Pr/Ph ratio of around 1, a sterane distribution of C₂₇>C₂₈≈C₂₉ (Fig.1), C₃₀ steranes, and a δ¹³C of the saturate and aromatic fractions of around -32 ‰ observed in several of the oils (e.g. Nizhne-Tabaganskoe, Yuzhno-Tabaganskoe) indicate that these can be correlated to the marine Jurassic Bazhenov source rock. A number of other oils (e.g. Maloichskoe, Urmanskoe, Severo-Ostaninskoe) are characterised by a sterane distribution dominated by C₂₉ steranes (Fig. 1), C₂₈ triaromatic steranes, δ¹³C of around -29 ‰ and a more waxy appearance in the alkane distribution. These oils can be correlated to the more terrigenous influenced Jurassic Togur/Tyumen source. Both sources have contributed to oils in various Paleozoic reservoirs in both the western and the south-eastern area indicating significant contributions of Mesozoic oils to Paleozoic reservoirs.

Concurrently, several oils cannot be correlated directly to either Jurassic source (e.g. Eley-Igayskoe) and are significantly different with respect to their biomarker composition implying at least one other, possibly marine carbonate, source. Whether this source is Paleozoic in origin, or another Jurassic horizon is as yet unknown, although a carbonate source rock would have to be Paleozoic, as no Mesozoic carbonate source rocks are known to date. However, the presence of another source contributing to Palaeozoic reservoirs, in a basin that is dominated by Mesozoic source rocks, provides significant scope for future exploration in the West Siberian Basin.

REFERENCES

- Goncharov, I., 1988. Geochemistry of oils from West Siberia. 204p.
- Kontorovich, A.E., Moskvina, V.I., Bostrikov, O.I., Danilova, V.P., Fomin, A.N., Fomichev, A.S., Kostyreva, E.A., and Melenevsky, V.N., 1996. Main oil source formations of the West Siberian basin: *Petroleum Geoscience* 3, 343-358.
- Rudkevich, M.Ya., Ozeranskaya, L.S., Chistyakova, H.F., Kornev, V.A., and Maksimov, E.M., 1988. Petroleum-productive complexes of the West Siberian basin (*Neftegazonosnye komplekсы Zapadno-Sibirskogo basseyna*): Moscow, Nedra, 304p.
- Peters, K.E., Kontorovich, A.E., Huizinga, B.J., Moldowan, J.M., and Lee, C.Y., 1994. Multiple oil families in the West Siberian basin: *American Association of Petroleum Geologists Bulletin* 78, 893-909.